

A P P L I C A T I O N

for

UNITED STATES LETTERS PATENT

on

MOLECULES THAT HOME TO A SELECTED ORGAN *IN VIVO*

by

Erkki Ruoslahti

and

Renata Pasqualini

CERTIFICATE OF MAILING BY "EXPRESS MAIL"

"EXPRESS MAIL" MAILING LABEL NUMBER: EL157452249US

DATE OF DEPOSIT: January 12, 1999

Sheets of Drawings: None

Docket No.: P-LJ 3430

I HEREBY CERTIFY THAT THIS PAPER OR FEE IS BEING  
DEPOSITED WITH THE UNITED STATES POSTAL SERVICE  
"EXPRESS MAIL POST OFFICE TO ADDRESSEE" SERVICE UNDER  
37 C.F.R. 1.10 ON THE DATE INDICATED ABOVE, AND IS  
ADDRESSED TO: THE ASSISTANT COMMISSIONER FOR PATENTS  
WASHINGTON, D.C. 20231.

Stephanie Fleischman

(TYPED OR PRINTED NAME OF PERSON MAILING PAPER OR FEE)

(SIGNATURE OF PERSON MAILING PAPER OR FEE)

Attorneys


CAMPBELL AND FLORES

4370 La Jolla Village Drive, Suite 700

San Diego, California 92122

09223866 "011299

## MOLECULES THAT HOME TO A SELECTED ORGAN IN VIVO

 This invention was made with government support under CA 42507, CA 62042 and Cancer Center Support Grant CA 30199 awarded by the National Institutes of Health.

5 The government has certain rights in this invention.

### BACKGROUND OF THE INVENTION

#### FIELD OF THE INVENTION

The present invention relates generally to the fields of molecular medicine and drug delivery and, more  
10 specifically, to molecules that home to a specific organ.

#### BACKGROUND INFORMATION

Although the effect of a particular pathology often is manifest throughout the body of the afflicted person, generally, the underlying pathology may affect  
15 only a single organ or tissue. In many cases, drugs are the treatment of choice for a patient suffering a particular disease. It is rare, however, that a drug will target only the diseased tissue or organ. More commonly, drug treatment results in undesirable side  
20 effects due, for example, to generalized toxic effects throughout the patient's body. The nausea, loss of hair and drop in blood count that occur as a result of treating of a cancer patient with chemotherapeutic agents are examples of the undesirable side effects that can  
25 occur due to drug treatment.

The undesirable side effects that can occur when drugs are used to treat a disease most often are due to the inability of the drug to specifically target the diseased organ or tissue. For example, a cancer  
30 chemotherapeutic agent that targets rapidly proliferating cells would be useful to kill rapidly dividing cancer

09223866-011299

cells. However, such an agent also kills normal  
proliferating hematopoietic and epithelial cells. Thus,  
the dose of such a drug that can be administered to a  
patient is limited due to its toxic effect on normal  
5 cells.

Efforts have been made to increase the target  
specificity of various drugs. In some cases, a  
particular cell type present in a diseased tissue or  
organ may express a unique cell surface marker. In such  
10 a case, an antibody can be raised against the unique cell  
surface marker and a drug can be linked to antibody.  
Upon administration of the drug/antibody complex to the  
patient, the binding of the antibody to the cell surface  
marker results in the delivery of a relatively high  
15 concentration of the drug to the diseased tissue or  
organ. Similar methods can be used where a particular  
cell type in the diseased organ expresses a unique cell  
surface receptor or a ligand for a particular receptor.  
In these cases, the drug can be linked to the specific  
20 ligand or to the receptor, respectively, thus providing a  
means to deliver a relatively high concentration of the  
drug to the diseased organ.

While linking a drug to a molecule that homes  
to a particular cell type present in a diseased organ or  
25 tissue provides significant advantages for treatment over  
the use of a drug, alone, use of this method is severely  
limited. In particular, very few cell type specific  
antibodies have been described and it can be difficult  
and time consuming to attempt to obtain an antibody that  
30 targets an organ in a particular patient suffering a  
pathology. Furthermore, few cell type specific surface  
markers have been described. Even where such markers  
have been described, the cells expressing the markers can  
be distributed among various tissues or organs, thereby  
35 limiting their usefulness as targets. Thus, it is

09228855-011299

important to identify specific target cell markers that are expressed in only one or a few tissues or organs and to identify molecules that specifically interact with such markers.

5                Various cell types can express unique markers and, therefore, provide potential targets for organ homing molecules. Endothelial cells, for example, which line the internal surfaces of blood vessels, can have distinct morphologies and biochemical markers in  
10 different tissues. The blood vessels of the lymphatic system, for example, express various adhesion proteins that serve to guide lymphocyte homing. For example, endothelial cells present in lymph nodes express a cell surface marker that is a ligand for L-selectin and  
15 endothelial cells in Peyer's patch venules express a ligand for the  $\alpha_4\beta_7$  integrin. These ligands are involved in specific lymphocyte homing to their respective lymphoid organs. Thus, linking a drug to L-selectin or to the  $\alpha_4\beta_7$  integrin may provide a means for targeting the  
20 drug to diseased lymph nodes or Peyer's patches, respectively, provided that these molecules do not bind to similar ligands present in a significant number of other organs.

25                Although the homing molecules present in the blood vessels of non-lymphoid tissues have not been clearly defined, the ability of lymphocytes to return to the organ in which they were first stimulated indicates that organ-specific endothelial markers exist. Similarly, the homing or metastasis of particular types  
30 of tumor cells to specific organs provides further evidence that organ-specific markers exist. However, there remains a need to identify other organ-specific cell markers and the molecules that bind to them.

09228866 "011299

Methods are now available for producing large populations of molecules. In addition, methods are available for screening libraries of molecules to identify those of interest. For example, phage peptide display libraries can be used to express large numbers of peptides that can be screened *in vitro* with a particular target molecule or a cell of interest in order to identify peptides that specifically bind the target molecule or the cell. Screening of such phage display libraries has been used, for example, to identify ligands that specifically bind various antibodies and cell surface receptors.

Screening of a phage display library generally involves *in vitro* panning of the library using a purified target molecule. Phage that bind the target molecule can be recovered, individual phage can be cloned and the peptide expressed by a cloned phage can be determined. Such a peptide can be useful for delivery of a drug linked to the peptide to cells expressing the target molecule.

Unfortunately, very few target molecules that are expressed by only one or a few cell types have been identified. Furthermore, even where such a target molecule is known, it is uncertain whether a peptide that specifically binds the target molecule, as determined using an *in vitro* panning method, will bind to the target molecule *in vivo*. As a result, the identification of a peptide from a phage display library using an *in vitro* panning method essentially represents only a starting point for determining whether the identified peptide can be useful for an *in vivo* procedure. Thus, a need exists to develop *in vivo* methods for screening large numbers of molecules such as peptides in order to identify those that can home to one or more selected organs. The

09228866.011299

present invention satisfies this need and provides related advantages as well.

### SUMMARY OF THE INVENTION

5 The present invention provides peptides that selectively home to a selected organ. For example, the invention provides brain homing peptides having the motif, SRL (serine-arginine-lysine), such as the peptide CLSSRLDAC (SEQ ID NO: 3), or the motif VLR, such as the peptide WRCVLREGPAGGCAWFNRHRL (SEQ ID NO: 16). In  
10 addition, the invention provides kidney homing peptides such as the peptide CLPVASC (SEQ ID NO: 21). An organ homing molecule of the invention is useful, for example, for targeting a moiety such as a drug to the selected organ or for identifying the presence of a target  
15 molecule in a sample.

In addition, the invention provides methods of identifying a target molecule by detecting selective binding of the target molecule to a peptide of the invention. For example, a peptide that selectively homes  
20 to a selected organ can be attached to a solid matrix for use in affinity chromatography. A sample of the organ can be obtained and passed over the affinity matrix under conditions that allow specific binding of the target molecule, which then can be collected and identified  
25 using well known biochemical methods. The target molecule can be useful, for example, for raising an antibody specific for the target molecule.

### DETAILED DESCRIPTION OF THE INVENTION

30 The present invention provides organ homing peptides, including brain homing peptides such as CNSRLHLRC (SEQ ID NO: 1), CENWWGDVC (SEQ ID NO: 2), WRCVLREGPAGGCAWFNRHRL (SEQ ID NO: 16) and others as shown

09228866-011299

in Table 1, and kidney homing peptides such as CLPVASC (SEQ ID NO: 21), CGAREMC (SEQ ID NO: 22) and others as shown in Table 2. Molecules that home specifically to one or a few selected organs were identified by screening a library using *in vivo* panning. The identified molecules are useful, for example, for targeting a desired moiety such as a drug, a toxin or a detectable label, which can be linked to the molecule, to the selected organ. *In vivo* panning provides a direct means for identifying molecules that specifically home to a selected organ and, therefore, provides a significant advantage over previous methods, which require that a molecule identified using an *in vitro* screening method subsequently be examined to determine whether it maintains its specificity *in vivo*.

As used herein, the term "library" means a collection of molecules. A library can contain a few or a large number of different molecules, varying from about ten molecules to several billion molecules or more. If desired, a molecule can be linked to a tag, which can facilitate recovery or identification of the molecule.

As used herein, the term "molecule" is used broadly to mean an organic chemical such as a drug; a peptide, including a variant or modified peptide or peptide-like molecules such as a peptidomimetic or peptoid; or a protein such as an antibody or a growth factor receptor or a fragment thereof such as an Fv, Fd or Fab fragment of an antibody, which contains a binding domain. For convenience, the term "peptide" is used broadly herein to mean peptides, proteins, fragments of proteins, peptoids, peptidomimetics and the like. A molecule can be a non-naturally occurring molecule, which does not occur in nature, but is produced as a result of *in vitro* methods, or can be a naturally occurring

09228866 "011299

molecule such as a protein or fragment thereof expressed from a cDNA library.

Methods for preparing libraries containing diverse populations of various types of molecules are well known in the art and commercially available (see, for example, Ecker and Crooke, Biotechnology 13:351-360 (1995), and Blondelle et al., Trends Anal. Chem. 14:83-92 (1995), and the references cited therein, each of which is incorporated herein by reference; see, also, Goodman and Ro, Peptidomimetics for Drug Design, in "Burger's Medicinal Chemistry and Drug Discovery" Vol. 1 (ed. M.E. Wolff; John Wiley & Sons 1995), pages 803-861, and Gordon et al., J. Med. Chem. 37:1385-1401 (1994), each of which is incorporated herein by reference). Where a molecule is a peptide, protein or fragment thereof, the molecule can be produced *in vitro* directly or can be expressed from a nucleic acid, which is produced *in vitro*. Methods of synthetic peptide and nucleic acid chemistry are well known in the art.

A library of molecules also can be produced, for example, by constructing a cDNA expression library from mRNA collected from a cell, tissue, organ or organism of interest. Methods for producing such libraries are well known in the art (see, for example, Sambrook et al., Molecular Cloning: A laboratory manual (Cold Spring Harbor Laboratory Press 1989), which is incorporated herein by reference). Preferably, the peptide encoded by the cDNA is expressed on the surface of a cell or a virus containing the cDNA. For example, cDNA can be cloned into a phage vector such as fuse 5 (see Example I), wherein, upon expression, the encoded peptide is expressed as a fusion protein on the surface of the phage.

As disclosed herein, *in vivo* panning comprises administering a library to a subject, collecting a selected organ and identifying an organ homing molecule. An organ homing molecule can be identified using various methods well known in the art. Generally, the presence of an organ homing molecule in a collected organ is identified based on one or more characteristics common to the molecules present in the library, then the structure of a particular organ homing molecule is identified. For example, a highly sensitive detection method such as mass spectrometry, either alone or in combination with a method such as gas chromatography, can be used to identify organ homing molecules in a selected organ. Thus, where a library comprises diverse molecules based generally on the structure of an organic molecule such as a drug, an organ homing molecule can be identified by determining the presence of a parent peak for the particular molecule.

If desired, the selected organ can be processed using a method such as HPLC, which can provide a fraction enriched in molecules having a defined range of molecular weights or polar or nonpolar characteristics or the like. Conditions for HPLC will depend on the chemistry of the particular molecule and are well known to those skilled in the art. Similarly, methods for bulk removal of potentially interfering cellular materials such as DNA, RNA, proteins, lipids or carbohydrates are well known in the art as are methods for enriching a fraction containing an organic molecule using, for example, methods of selective extraction. For example, where a library comprises a population of diverse organic chemical molecules each linked to a unique oligonucleotide tag, such that the specific molecule is identified by determining the oligonucleotide sequence using PCR, genomic DNA can be removed from the sample of the collected organ in order to reduce the potential for

09228866-011299

5

10  
15  
20  
25

30

35

smaller number of an untagged organic chemical homing molecule such as a drug is identifiable because such molecules normally are absent from or present in only small numbers in the body. In such a case, a highly sensitive method such as mass spectrometry can be used to identify an organ homing molecule. The skilled artisan will recognize that the method of identifying a molecule will depend, in part, on the chemistry of the particular molecule.

10           The molecules of a library also can be present as a conjugate, which can facilitate recovery or identification of the molecule. As used herein, the term "conjugate" or "molecule conjugate" means a molecule of the library linked to a physical, chemical or biological tag such as a plastic microbead, an oligonucleotide or a bacteriophage, respectively. The tag can provide a means to identify or recover an organ homing molecule of the invention following *in vivo* panning. For example, a conjugate can be a molecule such as a peptide linked to a unique oligonucleotide tag (see, for example, Brenner and Lerner, Proc. Natl. Acad. Sci., USA 89:5381-5383 (1992), which is incorporated herein by reference). Upon homing to an organ, the particular peptide can be identified by performing PCR on a sample of the organ containing the conjugate under conditions that allow amplification of the oligonucleotide tag (see, for example, Erlich, PCR Technology: Principles and Applications for DNA Amplification (Stockton Press 1989), which is incorporated herein by reference). By determining the oligonucleotide sequence, the identity of the peptide can be determined.

          In addition, a tag can be a support. As used herein, the term "support" means a tag having a defined surface to which a molecule can be attached. For example, a support can be a biological tag such as a

Where a molecule is linked to a support, the conjugate comprises the molecule attached to the surface of the support, such that the part of the molecule suspected of being able to interact with a target in a cell in the subject is positioned so as be able to participate in the interaction. For example, where the molecule is suspected of being a  $\beta$  adrenergic agonist, the binding portion of the molecule attached to a support is positioned so it can interact with a  $\beta$  adrenergic receptor on a cell in the selected organ. Similarly, where the molecule is suspected of being a ligand for a growth factor receptor, the molecule is positioned on the support so that it can bind the receptor.

As exemplified herein, a peptide suspected of being able to home to a selected organ such as brain was expressed as the N-terminus of a fusion protein, wherein the C-terminus consisted of a phage coat protein. Upon expression of the fusion protein, the C-terminal coat protein linked the fusion protein to the surface of a phage such that the N-terminal peptide was in a position to interact with a target molecule in an organ. Thus, a conjugate was formed by the linking of a peptide molecule

to a phage, wherein the phage provided a biological support and the peptide molecule is linked as a fusion protein.

As used herein, the term "in vivo panning" means a method of screening a library by administering the library to a subject and identifying a molecule that selectively homes to one or a few selected organs. The term "administering to a subject", when used in referring to a library of molecules or a portion thereof, is used in its broadest sense to mean that the library is delivered to a selected organ. For example, a library can be administered to a subject by injecting the library into the circulation of the subject such that the molecules can pass through the selected organ; after an appropriate period of time, circulation is terminated by sacrificing the subject or by removing a sample of the organ (see Example I). Alternatively, a cannula can be inserted into a blood vessel in the subject, such that the library is administered by perfusion for an appropriate period of time, after which the library can be removed from the circulation through the cannula or the subject can be sacrificed or the organ can be sampled to terminate circulation. Similarly, a library can be shunted through one or a few organs by cannulation of the appropriate blood vessels in the subject. It is recognized that a library also can be administered to an isolated perfused organ. Such panning in an isolated perfused organ can be useful for identifying molecules that bind to the organ and, if desired, can be used as an initial screening of a library. For example, if a kidney homing molecule is desired, a library can be perfused through an isolated kidney, then the molecules that bound to the perfused kidney can be screened by *in vivo* panning to identify a kidney homing molecule.

00228866-011299

The *in vivo* panning method is exemplified herein by screening a phage peptide display library in mice and identifying specific peptides that selectively home to brain or to kidney (see Examples I and II).

5 However, phage libraries that display protein receptor molecules, including, for example, an antibody or an antigen binding fragment of an antibody such as an Fv, Fd or Fab fragment; a hormone receptor such as a growth factor receptor; or a cell adhesion receptor such as an integrin  
10 or a selectin also can be used to practice the invention. Variants of such molecules can be constructed using well known methods such as random, site-directed or codon based mutagenesis (see Huse, U.S. Patent No. 5,264,563, issued November 23, 1993, which is incorporated herein by  
15 reference). Thus, various types of phage display libraries can be screened using the disclosed *in vivo* panning method.

Phage display technology provides a means for expressing a diverse population of random or selectively  
20 randomized peptides. Various methods of phage display and methods for producing diverse populations of peptides are well known in the art. For example, Ladner et al. (U.S. Patent No. 5,223,409, issued June 29, 1993, which is incorporated herein by reference) describe methods for  
25 preparing diverse populations of binding domains on the surface of a phage. In particular, Ladner et al. describe phage vectors useful for producing a phage display library, as well as methods for selecting potential binding domains and producing randomly or  
30 selectively mutated binding domains.

Similarly, Smith and Scott (Meth. Enzymol. 217:228-257 (1993); see, also, Scott and Smith, Science 249: 386-390 (1990), each of which is incorporated herein by reference) describe methods of producing phage peptide  
35 display libraries, including vectors and methods of

00228888-011299

diversifying the population of peptides that are expressed (see, also, Huse, WO 91/07141 and WO 91/07149, each of which is incorporated herein by reference; see, also, Example I). Phage display technology can be particularly powerful when used, for example, with a codon based mutagenesis method, which can be used to produce random peptides or randomly or desirably biased peptides (Huse, U.S. Patent No. 5,264,563, *supra*, 1993). These or other well known methods can be used to produce a phage display library, which can be subjected to the *in vivo* panning method of the invention in order to identify a peptide that homes to one or a few selected organs.

In addition, to screening a phage display library, *in vivo* panning can be used to screen various other types of libraries, including, for example, an RNA or cDNA library or a chemical library. If desired, the organ homing molecule can be in the form of a conjugate, which can facilitate recovery of the molecule from a selected organ or identification of the molecule in the organ. For example, where the molecules of a tagged chemical library are screened, the tag can be a moiety such as biotin, which can be linked directly to the molecule or can be linked to a support containing the molecules. Biotin provides a means to recover the molecule from a selected organ using an avidin or streptavidin affinity matrix. In addition, a molecule or a support containing a molecule can be linked to a hapten such as 4-ethoxy-methylene-2-phenyl-2-oxazoline-5-one (phOx), which can be bound by an anti-phOx antibody linked to a magnetic bead as a means to recover the molecule. Methods for purifying biotin or phOx labeled conjugates are known in the art and the materials for performing these procedures are commercially available (eg., Invitrogen; La Jolla CA; and Promega Corp.; Madison WI). In the case where a phage library is screened, the

phage can be recovered using methods as disclosed in Example I.

*In vivo* panning provides a method for directly identifying molecules that can home to one or a few selected organs. As used herein, the term "home" or "selectively home" means that a particular molecule binds relatively specifically to a target molecule present in one or few selected organs following administration to a subject. In general, selective homing is characterized, in part, by detecting at least a 2-fold (2x) greater specific binding of the molecule to the selected organ as compared to a control organ. It should be recognized that, in some cases, a molecule can localize non-specifically to an organ. Non-specific localization can be distinguished from homing by performing competition experiments such as those described in Examples II.C. and II.D.

Selective homing can be demonstrated by determining the specificity of an organ homing molecule for the selected organ as compared to a control organ. For example, a ratio of brain:kidney homing of up to 9:1 was observed for brain homing peptides (i.e., 9x greater binding to the selected organ as compared to the control organ; see Example II).

Selective homing also can be demonstrated by showing that molecules that home to a selected organ, as identified by one round of *in vivo* panning, are enriched for organ homing molecules in a subsequent round of *in vivo* panning. For example, phage expressing peptides that selectively home to brain were isolated by *in vivo* panning, then were subjected to additional rounds of *in vivo* panning. As demonstrated in Example II, phage recovered from brain after a first round of screening

showed an 8x enrichment in homing to brain as compared to kidney following a second round of screening and a 13x enrichment following a third round of *in vivo* panning.

When a peptide that selectively homes to brain was linked  
5 to a moiety, i.e., a red blood cell (RBC), the peptide/RBC complex selectively homed to brain (see Example II.D.).

Due to the conserved nature of cellular  
receptors and of ligands that bind a particular receptor,  
10 the skilled artisan would recognize that an organ homing molecule identified using, for example, *in vivo* panning in a mouse also would bind to the corresponding target molecule in the selected organ of a human or other species. For example, an RGD-containing peptide that can  
15 specifically bind to an integrin expressed by a cell in a human subject also can bind integrins expressed in a variety of species, including integrins expressed in mammalian cells such as murine and bovine cells as well as in cells of more evolutionarily distant species such  
20 as *Drosophila*. The ability of an organ homing molecule identified using *in vivo* panning in an experimental animal such as a mouse readily can be examined for the ability to bind to the corresponding organ in a human subject by demonstrating, for example, that the molecule  
25 also can bind specifically *in vitro* to a sample of the selected organ obtained from a human subject. Thus, routine methods can be used to confirm that an organ homing molecule identified using *in vivo* panning in an experimental animal also can bind an organ-specific  
30 target molecule in a human subject.

The steps of administering the library to the subject, collecting a selected organ and identifying the molecules that home to the selected organ, comprise a single round of *in vivo* panning. Although not required,

00228866-011299

one or more additional rounds of *in vivo* panning generally are performed. Where an additional round of *in vivo* panning is performed, the molecules recovered from the selected organ in the previous round are administered to a subject, which can be the same subject used in the previous round, where only a part of the selected organ was collected.

By performing a second round of *in vivo* panning, the relative binding selectivity of the molecule conjugates recovered from the first round can be determined by administering the identified molecule conjugates to a subject, collecting the selected organ and a control organ, and comparing the molecules recovered from the selected organ with those recovered from a control organ. In addition, a second round of *in vivo* panning can indicate whether the molecules identified from the initially selected organ also can home to additional organs, thus defining a family of selected organs. Additional rounds of panning can be performed as desired.

Ideally, no molecules are recovered from a control organ following a second or subsequent round of *in vivo* panning. Generally, however, a proportion of the molecules also will be present in a control organ. In this case, the ratio of molecules in the selected organ as compared to the control organ (selected:control) can be determined. As described above, for example, phage that homed to brain following a first round of *in vivo* panning demonstrated a 13x enrichment in homing to the selected organ as compared to the control organ, kidney, following two additional rounds of panning (Example II).

Additional rounds of *in vivo* panning can be used to determine whether a particular molecule homes only to the selected organ or can recognize a target on the selected organ that also is expressed in one or more other organs or is sufficiently similar to the target in the originally selected organ. Where a molecule is found to direct homing to organs in addition to the originally selected organ, the organs are considered to constitute a family of selected organs. Using the method of *in vivo* panning, molecules that home to only a single selected organ and molecules that home to a family of selected organs can be defined. Such identification is expedited by collecting various organs during subsequent rounds of *in vivo* panning.

As used herein, the term "selected organ" is used in its broadest sense to mean an organ to which a molecule selectively homes. In addition, the term "organ" is used broadly to mean organ, tissue or cell type, including a cancer cell, in which case the selected organ can be a tumor such as a primary tumor or a metastatic lesion. In general, a selected organ contains a cell that expresses a particular target molecule such as a cell surface receptor to which an organ homing molecule can bind. By performing at least two rounds of *in vivo* panning, the selectivity of homing of the molecule to the selected organ can be determined (see Example II). As discussed above, however, in some cases an organ homing molecule can selectively home to more than one selected organ, in which case the molecule is considered to be able to selectively home to a family of selected organs.

The term "control organ" is used to mean an organ other than a selected organ. A control organ is characterized by the inability of an organ homing

molecule to home to the control organ. A control organ is useful for identifying non-specific binding of a molecule. A control organ can be collected, for example, to identify non-specific binding of the molecule or to  
5 determine the selectivity of homing of the molecule (see Examples I and II). In addition, non-specific binding can be identified by administering, for example, a control molecule, which is known not to home to an organ but is chemically similar to a potential organ homing  
10 molecule. Alternatively, where the molecules are administered as a conjugate comprising the molecules of the library linked to a support, administration of the supports, alone, can be used to identify non-specific binding. For example, a phage that expresses the gene  
15 III protein, alone, but that does not contain a peptide fusion protein, can be screened by *in vivo* panning to determine the level of non-specific binding of the phage support.

In some cases it can take several rounds of *in*  
20 *vivo* panning to identify a control organ, since a molecule that selectively homes to an originally selected organ also may have the ability to selectively home to additional organs, thus defining a family of selected organs. The ability of a molecule to home to one or to a  
25 family of selected organs allows for the preparation of panels of organ homing molecules, wherein individual molecules variously home to one selected organ or to any of a family of selected organs with variable selectivity.

In general, a library of molecules, which  
30 contains a diverse population of random or selectively randomized molecules of interest, is prepared, then administered to a subject. At a selected time after administration, the subject is sacrificed and a selected organ or part of the organ is collected such that the  
35 molecules present in the selected organ can be

00000000.011299

5

10

25

30

vivo or whether the binding peptide or the receptor are unique to a specific organ in the body. Furthermore, the *in vitro* methods are performed using defined, well-characterized target molecules in an artificial system.

- 5 For example, Goodson et al. utilized cells expressing a recombinant urokinase receptor. Thus, the *in vitro* methods require prior knowledge of the target molecule and yield little if any information regarding *in vivo* utility. In contrast, the *in vivo* panning method
- 10 disclosed herein requires no prior knowledge or availability of a target molecule. Thus, *in vivo* panning provides a significant advantage over previous methods by identifying molecules that selectively home *in vivo* to a target molecule present in one or a few organs.

- 15 Brain and kidney were selected as target organs to identify phage expressing peptides that selectively home to these selected organs because the blood vessels in these organs have unique characteristics. For example, the blood vessels in the brain form the
- 20 "blood-brain barrier" and express at least one specific antigen (Schlosshauer and Herzog, J. Cell Biol. 110:1261-1274 (1990)). Using *in vivo* panning, phage expressing various peptides that selectively homed to brain or to kidney, but not to both organs, were identified (see
- 25 Example II). Since the *in vivo* panning was terminated about one to four minutes after intravenous administration of the phage to the mice, the majority of phage likely remained in the vasculature, indicating the displayed peptides bound to endothelial cell surface
- 30 markers that are expressed in an organ-specific manner (see, for example, Springer, Cell 76:301-314 (1994)). Organ-specific homing of tumor cells and of lymphocytes indicates that the vascular beds in various tissues express unique target molecules. For example, the

00228866-011299

ability of lymphocytes to return to the organ in which they were first stimulated (Salmi et al., Proc. Natl. Acad. Sci., USA 89:11436-11440 (1992)) and the ability of particular types of tumor cells to home or metastasize to specific organs (Fidler and Hart, Science 217:998-1003 (1982)) indicate that organ-specific targets are expressed in a subject.

The identification of brain homing and kidney homing peptides demonstrates that organ-specific target molecules exist and indicates that the disclosed method of *in vivo* panning can be used to identify molecules that selectively home to other organs. For example, tumor vasculature, which undergoes active angiogenesis and contains specific markers (Brooks et al., Cell 79:1157-1164 (1994)), is a particularly attractive target. The identification of molecules that home to tumor vasculature would provide a means for directing a therapeutic agent directly to a tumor while sparing normal tissues.

Phage peptide display libraries were constructed essentially as described Smith and Scott (*supra*, 1993; see, also, Koivunen et al., Biotechnology 13:265-270 (1995), which is incorporated herein by reference). Oligonucleotides encoding peptides having substantially random amino acid sequences were synthesized based on an "NNK" codon, wherein "N" is A, T, C or G and "K" is G or T. "NNK" encodes 32 triplets, which encode the twenty amino acids and an amber STOP codon (Scott and Smith, *supra*, 1990). At least one codon encoding cysteine also was included in each oligonucleotide so that cyclic peptides could be formed through disulfide linkages (see Example I). The oligonucleotides were inserted in frame with the sequence encoding the gene III protein (gIII) in the vector fuse 5

09228866 "011299

such that a peptide-gIII fusion protein can be expressed. Following expression, the fusion protein is expressed on the surface of the phage containing the vector (Smith and Scott, *supra*, 1993; Koivunen et al., *supra*, 1994b).

5                   Remarkably, following *in vivo* panning, the phage that selectively homed to brain or to kidney displayed only a few different peptide sequences. In some cases, peptides having the same amino acid sequence were encoded by phage having different oligonucleotide  
10 sequences encoding the peptide. Furthermore, a family of brain homing peptides was identified, wherein each peptide in the family contained the common amino acid motif, SRL (serine-arginine-leucine), but different flanking amino acid sequences (see Table 1; SEQ ID NOS:  
15 1, 3 and 5). In addition, two different peptides displayed the motif VLR (valine-leucine-arginine; see Table 1, SEQ ID NOS: 4 and 16). These results demonstrate that it is the peptide displayed by the phage, rather than some incidental mutant property of the  
20 phage, that directs homing to the selected organ.

The sequences of the brain and kidney homing motifs do not reveal any significant similarities with known ligands for endothelial cell receptors nor do they resemble any sequences listed in various data banks.  
25 However, some of the brain-homing motifs share a similarity with integrin-binding sequences. For example, one brain homing peptide contained an RLD sequence, which is recognized by certain integrins (Altieri et al., J. Biol. Chem. 265:12119-12122 (1990); Koivunen et al.,  
30 *supra*, 1994a), and the DXXR (SEQ ID NO: 44) motif in another peptide resembles the RGD, DGR, and NGR motifs that bind to certain integrins (Ruoslahti, J. Clin. Invest. 87:1-5 (1991); Koivunen et al., *supra*, 1994a).

Selective homing of a molecule such as a peptide or protein to a selected organ can be due to specific recognition by the peptide of a particular cell target molecule such as a cell surface receptor present on a cell in the organ. Selectivity of homing is dependent on the particular target molecule being expressed on only one or a few different cell types, such that the molecule homes to only one or a few organs. As discussed above, brain and kidney homing peptides likely are recognizing endothelial cell surface markers in the blood vessels present in these organs. However, most different cell types, particularly cell types that are unique to an organ, can express unique target molecules. Thus, in organs such as liver, spleen or lymph node, where blood circulates through sinusoids formed by the cells specific for the organ, *in vivo* panning can be useful for identifying molecules that home to the particular organ. For example, the sinusoids in liver are formed, in part, by hepatocytes, which are unique to liver. Using the methods disclosed herein, *in vivo* panning can be used to identify molecules that selectively home to hepatocytes and specific homing can be demonstrated by performing the appropriate competition experiments.

An organ homing molecule such as a brain homing (Table 1) or kidney homing (Table 2) peptide can be used to direct a moiety to a selected organ by linking the moiety to the molecule. As used herein, the term "moiety" is used broadly to mean an agent linked to the organ homing molecule. A moiety can be, for example, a detectable label such a radiolabel; a toxin such as ricin; or a drug such as a chemotherapeutic agent. Various moieties and methods for linking a moiety to a molecule are well known in the art and commercially available (see, for example, Harlow and Lane, Antibodies:

A laboratory manual (Cold Spring Harbor Laboratory Press 1988), which is incorporated herein by reference).

Linking of a moiety to an organ homing molecule for the purpose of directing homing of the moiety to the selected organ is exemplified by the linking of a brain homing peptide to a RBC, wherein the peptide directed homing of the RBC to brain (see Example II.D.). These results indicate that an organ homing molecule of the invention can be linked to other cell types in order to direct the cell type to a selected organ. For example, a tumor homing molecule identified by *in vivo* panning can be linked to a white blood cell (WBC) such as a cytotoxic T cell or a killer cell, wherein upon administration of the tumor homing molecule/WBC complex, the molecule directs homing of the WBC to the tumor, where the WBC can exert its effector function.

An organ homing molecule also can be linked to a moiety that is detectable external to the subject in order to perform an *in vivo* diagnostic imaging study. For example, *in vivo* imaging using a detectably labeled brain homing peptide can identify a region in the brain where circulation is occluded. For such studies, a gamma ray emitting radionuclide such as indium-111 or technitium-99 can be linked to a brain homing molecule and, following administration to a subject, can be detected using a solid scintillation detector. Alternatively, a positron emitting radionuclide such as carbon-11 or a paramagnetic spin label such as carbon-13 can be linked to the molecule and, following administration to a subject, the localization of the moiety/molecule can be detected using positron emission transaxial tomography or magnetic resonance imaging, respectively.

Such *in vivo* imaging methods also can be used to identify the presence of cancer in a subject by linking an appropriate moiety to a tumor homing molecule, which can recognize a unique target expressed by the tumor cells or by the blood vessels formed by angiogenesis in the tumor. Such methods can identify a primary tumor as well as a metastatic lesion, which may not be detectable using other methods. Having identified the presence of such cancer, the tumor homing molecule can be linked to a toxin or to a chemotherapeutic agent in order to direct the moiety to the tumor. Such a method can allow selective killing of the tumor, while substantially sparing normal tissues.

When administered to a subject, the molecule/moiety complex is administered as a pharmaceutical composition containing, for example, the complex and a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers are well known in the art and include, for example, aqueous solutions such as water or physiologically buffered saline or other solvents or vehicles such as glycols, glycerol, oils such as olive oil or injectable organic esters.

A pharmaceutically acceptable carrier can contain physiologically acceptable compounds that act, for example, to stabilize or to increase the absorption of the complex. Such physiologically acceptable compounds include, for example, carbohydrates, such as glucose, sucrose or dextrans, antioxidants, such as ascorbic acid or glutathione, chelating agents, low molecular weight proteins or other stabilizers or excipients. One skilled in the art would know that the choice of a pharmaceutically acceptable carrier, including a physiologically acceptable compound, depends, for example, on the route of administration of the

09220806 011299

[illegible][illegible][illegible][illegible]

particularly useful when administered orally because they can be degraded in the digestive tract. However, methods for chemically modifying peptides to render them less susceptible to degradation by endogenous proteases or more absorbable through the alimentary tract are well known (see, for example, Blondelle et al., *supra*, 1995; Ecker and Crooke, *supra*, 1995; Goodman and Ro, *supra*, 1995). Such methods can be performed on peptides identified by *in vivo* panning. In addition, methods for preparing libraries of peptide analogs such as peptides containing D-amino acids; peptidomimetics consisting of organic molecules that mimic the structure of peptide; or peptoids such as vinylogous peptoids, are known in the art and can be used to identify molecules that home to a selected organ and are stable for oral administration.

Organ homing molecules obtained using the methods disclosed herein also can be useful for identifying a target molecule such as a cell surface receptor or a ligand for a receptor, which is recognized by the organ homing peptide. For example, an organ homing peptide can be linked to a solid support such as a chromatography matrix. The linked peptide then can be used for affinity chromatography by passing an appropriately processed sample of a selected organ over the column in order to bind a particular target molecule. The bound target molecule can be eluted from the column and can be identified and characterized by various well known methods. In addition, an organ homing peptide can be linked to a detectable moiety such as a radionuclide, a fluorescent molecule, an enzyme or biotin and can be used, for example, to screen a sample in order to detect the target molecule during various steps of purification of the target molecule.

The present invention also provides peptides that selectively home to a selected organ. For example,

Cysteine residues were included in the peptides such that cyclization of the peptides could be effected. However, in some cases one or both of the cysteine residues in a peptide can be deleted without significantly affecting the organ homing activity of a peptide of the invention. Thus, a peptide having the sequence LSSRLDA (SEQ ID NO: 19; compare SEQ ID NO: 3) also can be a brain homing peptide. Similarly, the amino acid residues N-terminal and C-terminal to the first and last cysteine residues, respectively, in a peptide such as WRCVLREGPAGGCAWFNRHRL (SEQ ID NO: 16) can be dispensable without substantially altering brain homing activity of the peptide. Thus, a peptide having the sequence VLREGPAGG (SEQ ID NO: 20) also can be useful as a brain homing peptide. Methods for determining the necessity of a cysteine residue or of amino acid residues N-terminal or C-terminal to a cysteine residue for organ homing activity of a peptide of the invention are routine and well known in the art.

30           An organ homing peptide of the invention is useful, for example, for targeting a desired moiety to the selected organ as discussed above. In addition, a peptide of the invention can be used to identify the presence of a target molecule in a sample. As used  
35 herein, the term "sample" is used in its broadest sense

5 If desired, a sample can be processed, for example, by homogenization, which can be an initial step for isolating the target molecule to which an organ homing molecule binds.

An organ homing peptide of the invention such as a brain homing peptide can be used to identify the target molecule expressed in brain. For example, a brain homing peptide can be attached to a matrix such as a chromatography matrix to produce a peptide affinity matrix. A homogenized sample of brain can be applied to the peptide-affinity matrix under conditions that allow specific binding of the brain homing peptide to the target molecule (see, for example, Deutscher, Meth. Enzymol., Guide to Protein Purification (Academic Press, Inc., ed. M.P. Deutscher, 1990), Vol. 182, which is incorporated herein by reference; see, for example, pages 357-379). Unbound and non-specifically bound material can be removed and the specifically bound brain-derived target molecule can be isolated in substantially purified form.

25           The following examples are intended to  
illustrate but not limit the present invention.

## IN VIVO PANNING

This example demonstrates methods for preparing  
30 a phage library and screening the library using *in vivo*  
panning to identify phage expressing peptides that home  
to a selected organ.

### A. Preparation of phage libraries:

Phage display libraries were constructed using the fuse 5 vector as described by Koivunen et al. (*supra*, 1995; see, also, Koivunen et al. Meth. Enzymol.

5 245:346-369 (1994b), which is incorporated herein by reference). Six libraries encoding peptides designated CX<sub>5</sub>C (SEQ ID NO: 36), CX<sub>6</sub>C (SEQ ID NO: 37), CX<sub>7</sub>C (SEQ ID NO: 38), CX<sub>9</sub> (SEQ ID NO: 39), X<sub>2</sub>CX<sub>14</sub>CX<sub>2</sub> (SEQ ID NO: 40) and X<sub>2</sub>CX<sub>18</sub> (SEQ ID NO: 41) were prepared, where "C" indicates  
 10 cysteine and "X<sub>N</sub>" indicates the given number of individually selected amino acids. These libraries can display cyclic peptides when at least two cysteine residues are present in the peptide.

Oligonucleotides were constructed such that "C"  
 15 was encoded by the codon TGT and "X<sub>N</sub>" was encoded by NNK, where "N" is equal molar mixtures of A, C, G and T, and where "K" is equal molar mixtures of G and T. Thus, the peptide represented by CX<sub>5</sub>C (SEQ ID NO: 36) can be represented by an oligonucleotide having the sequence  
 20 TGT(NNK)<sub>5</sub>TGT (SEQ ID NO: 42). Oligonucleotides were made double stranded by 3 cycles of PCR amplification, purified and ligated to the nucleic acid encoding the gene III protein in the fuse 5 vector such that, upon  
 25 expression, the peptide is present as a fusion protein at the N-terminus of the gene III protein.

The vectors were transfected by electroporation into MC1061 cells. Bacteria were cultured for 24 hr in the presence of 20 µg/ml tetracycline, then phage were collected from the supernatant by precipitation twice  
 30 using polyethylene glycol. Each library contained about 5 x 10<sup>9</sup> to 5 x 10<sup>14</sup> transducing units (TU; individual recombinant phage).

09228866-011299

B. In vivo panning of phage:

A mixture of phage libraries containing  $1 \times 10^{14}$  TU was diluted in 200  $\mu$ l DMEM and injected into the tail vein of anesthetized BALB/c mice (2 month old females; Jackson Laboratories; Bar Harbor ME); Avertin (0.015 ml/g) was used as anesthetic. After 1-4 minutes, mice were snap frozen in liquid nitrogen. To recover the phage, carcasses were partially thawed at room temperature for 1 hr, organs were collected and weighed, then were ground in 1 ml DMEM-PI (DMEM containing protease inhibitors (PI); phenylmethylsulfonyl fluoride (PMSF; 1 mM), aprotinin (20  $\mu$ g/ml), leupeptin (1  $\mu$ g/ml)).

Organ samples were washed 3 times with ice cold DMEM-PI containing 1% bovine serum albumin (BSA), then directly incubated with 1 ml K91-kan bacteria for 1 hr. Ten ml NZY medium containing 0.2  $\mu$ g/ml tetracycline (NZY/tet) was added to the bacterial culture, the mixture was incubated in a 37°C shaker for 1 hr, then 200  $\mu$ l aliquots were plated in agar plates containing 40  $\mu$ g/ml tetracycline (tet/agar).

Individual colonies containing phage recovered from brain or kidney were grown for 16 hr in 5 ml NZY/tet. The bacterial cultures obtained from the individual colonies were pooled and the amplified eluates were injected into mice as described above for a second round of *in vivo* panning. A third round of panning also was performed. Phage DNA was purified from individual bacterial colonies obtained from the second and the third round of *in vivo* panning and the DNA sequences encoding the peptides expressed by selected phage were determined (see Koivunen et al., *supra*, 1994b).

## EXAMPLE II

CHARACTERIZATION OF PEPTIDES THAT HOME  
TO A SELECTED ORGAN

This example demonstrates that an organ homing  
5 peptide of the invention selectively homes to a selected  
organ and that an organ homing peptide identified by *in*  
*vivo* panning can be used to direct a moiety to a selected  
organ.

A. Brain is the selected organ

10 Three rounds of *in vivo* panning in mice were  
performed. Kidney was used as a control organ. Mice  
were injected with two different mixtures of phage  
libraries. The first mixture contained libraries  
encoding CX<sub>9</sub> (SEQ ID NO: 39), CX<sub>5</sub>C (SEQ ID NO: 36), CX<sub>6</sub>C  
15 (SEQ ID NO: 37) and CX<sub>7</sub>C (SEQ ID NO: 38) peptides  
(CX<sub>5-7</sub>/CX<sub>9</sub> mixture; SEQ ID NOS: 36-39). The second mixture  
contained libraries encoding X<sub>2</sub>CX<sub>14</sub>CX<sub>2</sub> (SEQ ID NO: 40) and  
X<sub>2</sub>CX<sub>18</sub> (SEQ ID NO: 41) peptides (X<sub>2</sub>CX<sub>18</sub>/X<sub>2</sub>CX<sub>14</sub>CX<sub>2</sub> mixture;  
SEQ ID NOS: 40 and 41).

20 The phage library mixtures were administered to  
mice via tail vein injection. Phage input was  $1 \times 10^{16}$  TU  
of the CX<sub>5-7</sub>/CX<sub>9</sub> (SEQ ID NOS: 36-39) mixture or  $1 \times 10^{14}$  TU  
of the X<sub>2</sub>CX<sub>18</sub>/X<sub>2</sub>CX<sub>14</sub>CX<sub>2</sub> (SEQ ID NOS: 40 and 41) mixture.  
Phage were recovered from the brains of the injected  
25 mice, the recovered phage were amplified *in vitro*, then a  
second and a third round of *in vivo* panning was  
performed. During the second and third rounds of  
panning, phage were recovered from brain and from kidney  
and the number of TU from each organ was compared. This  
30 comparison revealed that 6x more phage from the CX<sub>5-7</sub>/CX<sub>9</sub>  
(SEQ ID NOS: 36-39) mixture bound to brain than to kidney

00228866 011299

5

10

25

35

organ is due to the specific peptide expressed on the phage.

To determine the specificity of brain homing of the individual motifs identified, phage displaying the predominant motifs were amplified individually, diluted to the same input titer and administered to mice. Following administration, brain and kidney were removed and the number of TU of phage in each organ was determined. The enrichment ratio of phage recovered from the selected organ, brain, as compared to the control organ, kidney, revealed that phage displaying one of the four most recovered peptides, CNSRLHLRC (SEQ ID NO: 1), CENWWGDVC (SEQ ID NO: 2), WRCVLREGPAGGCAWFNRHRL (SEQ ID NO: 16), CLSSRLDAC (SEQ ID NO: 3), each selectively targeted the brain as compared to the kidney.

[illegible]

TABLE 1

## PEPTIDES FROM PHAGE RECOVERED FROM BRAIN

5	A.A. SEQUENCE (SEQ ID NO:)	# PHAGE DISPLAYING SAME A.A. SEQUENCE (% PHAGE)
<u>CX<sub>5-7</sub>C/CX<sub>9</sub> (36-39) library:</u>		
	CNSRLHLRC (1)	16 (21.9%)
10	CENWWGDVC (2)	14 (19.2%)
	CLSSRLDAC (3)	6 (8.2%)
	CVLRGGRC (4)	5 (6.8%)
	CNSRLQLRC (5)	4 (5.5%)
	CGVRLGC (6)	3 (4.1%)
15	CKDWGRIC (7)	2 (2.8%)
	CLDWGRIC (8)	2 (2.8%)
	CTRITESC (9)	2 (2.8%)
	CETLPAC (10)	1 (1.4%)
	CRTGTLFC (11)	1 (1.4%)
20	CGRSLDAC (12)	1 (1.4%)
	CRHWFDVVC (13)	1 (1.4%)
	CANAQSHC (14)	1 (1.4%)
	CGNPSYRC (15)	1 (1.4%)
<u>X<sub>2</sub>CX<sub>18</sub>/X<sub>2</sub>CX<sub>14</sub>CX<sub>2</sub> (SEO ID NOS: 40 and 41) library:</u>		
25	WRCVLREGPAGGCAWFNRHRL (16)	10 (13.7%)
	YPCGGEAVAGVSSVRTMCSE (17)	1 (1.4%)
	LNCDYQGTNPATSVSVPCTV (18)	1 (1.4%)

09228866-011299

Specifically, the ratio of selective homing (brain:kidney) was about 8 for CNSRLHLRC (SEQ ID NO: 1) and for CLSSRLDAC (SEQ ID NO: 3), 4 for CENWWGDVC (SEQ ID NO: 2) and 9 for WRCVLREGPAGGCAWFNRHRL (SEQ ID NO: 16).

- 5 Two control phage showed low binding to both brain and kidney. These results demonstrate that *in vivo* panning can be used to screen phage display libraries in order to identify phage expressing peptides that home to a selected organ.

#### 10 B. Kidney as the selected organ

- The same methodology used to isolate phage expressing brain homing peptides was used to isolate phage expressing peptides that home to kidney. In these experiments, brain was used as the control organ. A  
15 mixture of the CX<sub>5</sub>C (SEQ ID NO: 36) and the CX<sub>6</sub>C (SEQ ID NO: 37) libraries was administered as described above. Homing of phage to the kidney was obtained and an approximately 3x to 7x enrichment of phage homing to kidney was observed following a second round of *in vivo*  
20 panning.

- The amino acid sequences were determined for the inserts in 48 cloned phage that homed to kidney. The peptides expressed by these phage were represented by two predominant sequences, CLPVASC (SEQ ID NO: 21; 46% of the  
25 clones sequenced) and CGAREMC (SEQ ID NO: 22; 17% of the clones; see Table 2). In addition, the peptide CKGRSSAC (SEQ ID NO: 23) appeared three times and three other peptides were present twice each. Phage expressing a  
30 CLPVASC (SEQ ID NO: 21), CGAREMC (SEQ ID NO: 22) or CKGRSSAC (SEQ ID NO: 23) peptide each exhibited selective homing to kidney; the ratio of selective kidney:brain homing was 7 for CLPVASC (SEQ ID NO: 21), 3 for CGAREMC (SEQ ID NO: 22) and 2 for CKGRSSAC (SEQ ID NO: 23).

00228866-011299

TABLE 2

PEPTIDES FROM PHAGE RECOVERED FROM KIDNEY

5	A.A. SEQUENCE (SEQ ID NO:)		# PHAGE DISPLAYING SAME A.A. SEQUENCE (% PHAGE)
10	CLPVASC	(21)	22 (45.8%)
	CGAREMC	(22)	8 (16.7%)
	CKGRSSAC	(23)	3 (6.2%)
	CWARAQGC	(24)	2 (4.2%)
	CLGRSSVC	(25)	2 (4.2%)
15	CTSPGGSC	(26)	2 (4.2%)
	CMGRWRLC	(27)	1 (2.1%)
	CVGECGGC	(28)	1 (2.1%)
	CVAWLNC	(29)	1 (2.1%)
	CRRFQDC	(30)	1 (2.1%)
20	CLMGVHC	(31)	1 (2.1%)
	CKLLSGVC	(32)	1 (2.1%)
	CFVGHDLC	(33)	1 (2.1%)
	CRCLNVC	(34)	1 (2.1%)
	CKLMGEC	(35)	1 (2.1%)

These results demonstrate that the *in vivo* panning method is a generally applicable method for screening a phage library to identify phage expressing peptides that home to a selected organ. Database searches did not reveal any significant homology of the brain homing or kidney homing peptides to known ligands for endothelial cell receptors.

C. Peptide homing is specific:

In order to confirm the specificity of a peptide for directing homing to a selected organ, peptide competition experiments were performed. The cyclic peptide, CLSSRLDAC (SEQ ID NO: 3), which is one of the brain homing peptides (see Table 1) was synthesized (Immunodynamics; La Jolla CA) and purified by HPLC. The effect on the homing of phage expressing CLSSRLDAC (SEQ ID NO: 3), CENWWGDVC (SEQ ID NO: 2) or WRCVLREGPAGGCAWFNRHRL (SEQ ID NO: 16) to the brain was examined in order to determine whether co-administration of the synthetic peptide affected homing of the phage.

Phage expressing CLSSRLDAC (SEQ ID NO: 3), CENWWGDVC (SEQ ID NO: 2) or WRCVLREGPAGGCAWFNRHRL (SEQ ID NO: 16) were titrated to the same concentration and  $1 \times 10^8$  TU was injected into mice alone, or with 100  $\mu$ g purified synthetic cyclic CLSSRLDAC (SEQ ID NO: 3) peptide. The synthetic CLSSRLDAC (SEQ ID NO: 3) peptide inhibited the homing of phage expressing CLSSRLDAC (SEQ ID NO: 3) by about 60%. This result demonstrates that the homing of the phage to brain is specifically due to the expression on the phage of the CLSSRLDAC (SEQ ID NO: 3) peptide.

The synthetic CLSSRLDAC (SEQ ID NO: 3) peptide also inhibited homing of phage expressing the WRCVLREGPAGGCAWFNRHRL (SEQ ID NO: 16) peptide by about 60%, but did not affect homing of CENWWGDVC (SEQ ID NO: 2) phage. This result indicates that the CLSSRLDAC (SEQ ID NO: 3) and WRCVLREGPAGGCAWFNRHRL (SEQ ID NO: 16) peptides can recognize the same target molecule in brain, whereas the CENWWGDVC (SEQ ID NO: 2) peptide recognizes a different target.

09228866-011299

D. The brain homing peptide CLSSRLDAC (SEQ ID NO: 3)  
directs red blood cells to brain

The synthetic cyclic CLSSRLDAC (SEQ ID NO: 3)  
5 peptide (1 mg) was labeled with iodine-125 using the  
Bolton and Hunter reagent (Amersham; Arlington Heights  
IL). Labeled peptide was purified by reverse phase HPLC  
on Sep-Pak<sup>TM</sup> cartridges (Waters, Millipore Corp.; Milford  
PA) and the <sup>125</sup>I-peptide (200 µg) was coupled to 1 ml  
10 glutaraldehyde-stabilized sheep RBC (Sigma Chemical Co.;  
St. Louis MO) according to the manufacturer's  
instructions.

Fifty µl <sup>125</sup>I-peptide/RBC (200,000 cpm), alone  
or with 10 mM of unlabeled CLSSRLDAC (SEQ ID NO: 3),  
15 which is a brain homing peptide, or CVRLNSLAC (SEQ ID  
NO: 43), which has no brain homing activity, was injected  
into the tail vein. After 2 min, each mouse was perfused  
through the heart with 50 ml DMEM and the brain and  
kidney were removed and assayed for radioactivity.

20 Approximately twice as much of the  
CLSSRLDAC/RBC complex (SEQ ID NO: 3) homed to brain than  
to kidney. Coadministration of unlabeled CLSSRLDAC (SEQ  
ID NO: 3) with the complex essentially completely  
inhibited brain homing of the complex but had no effect  
25 on complex localizing to the kidney, indicating that  
localization of the complex in the kidney was  
non-specific. Coadministration of unlabeled CVRLNSLAC  
(SEQ ID NO: 43) had no effect on selective homing of the  
complex to brain and no effect on the non-specific  
30 localization of the complex in the kidney. These results  
demonstrate that an organ homing peptide identified using  
in vivo panning can be linked to a moiety such as a blood  
cell and selectively directs homing of the linked moiety  
to the selected organ.

0923886-01299

Although the invention has been described with reference to the disclosed examples, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly,  
5 the invention is limited only by the following claims.

09228866-011299  
662710-99882360